

# Solving Assignment Problem Using Heptagonal Fuzzy Number

<sup>1</sup>A. Deepa , <sup>2\*</sup>S. Santhi

<sup>1</sup>Research Scholar, Department of Mathematics,

Vels Institute of Science Technology and Advanced Studies, Pallavaram, Tamil Nadu, India.

Email: adeepa004@gmail.com

<sup>2</sup>Assistant Professor, Department of Mathematics,

Vels Institute of Science Technology and Advanced Studies, Pallavaram, Tamil Nadu, India.

\*Corresponding Author Email : Santhosh.mitha@gmail.com

**Received:** 2022 March 15; **Revised:** 2022 April 20; **Accepted:** 2022 May 10.

---

## Abstract

Now a days Assignment problems were widely applied in the field of manufacturing sectors, production planning ,etc to earn more profit with minimum inventory. Generally Fuzzy Numbers plays a vital role in many decision making problems. Here an optimal solution of the assignment problem had been derived from Hungarian method .The parameters were being given by using Heptagonal fuzzy numbers which are further decomposed into pentagonal fuzzy numbers. The fuzzy numbers were defuzzified into real numbers by using pascal's graded mean technique. Numerical example had been given to illustrate the solution procedure.

**Keywords:**Assignment problem, Hungarian Method ,Heptagonal fuzzy numbers, Pentagonal fuzzy numbers, Optimization Technique.

---

**Introduction:** An Assignment problem is one of the fundamental combinatorial problem in the branch of operation research. The main objective of the assignment problem is to assign number of resources to an equal number of activities

in order to minimize the total cost (or) to maximize the profit. Assignment Problem is widely applied in manufacturing sectors and many service systems. Many algorithms had been formulated to find the optimal solution .One of the best known

algorithm is Hungarian Algorithm. ASahaya Sudha and S.Karunambigai Solved the Transportation Problem Using Heptagonal Fuzzy Numbers[6]. R. Srinivasan, N. Karthikeyan, A. Jayaraja[1] proposed a ranking method for pentagonal fuzzy number in solving transportation problem. Shweta Singh, G.C. Dubey, Rajesh Shrivastava [2] made a Comparative Analysis of Assignment Problem. Here an assignment problem had been solved by Hungarian method using Heptagonal Fuzzy Number, further it had been decomposed into Pentagonal fuzzy number and analysed the results obtained.

## Preliminaries

### Fuzzy Set

Let  $X$  be a universal set. The Fuzzy Set  $\tilde{A}$  is defined by the membership function  $\mu_{\tilde{A}}(x)$  that maps the element of domain space into unit interval  $[0,1]$  where  $\mu_{\tilde{A}}$  is the degree of membership of  $x \in X$  in the set  $\tilde{A}$  and is denoted by  $\tilde{A} = \{x, \mu_{\tilde{A}}(x) : x \in X\}$ .

### Fuzzy Number

The Fuzzy Number  $\tilde{A}$  is a fuzzy set whose membership function  $\mu_{\tilde{A}}(x)$  satisfies the following conditions

- (i)  $\mu_{\tilde{A}}(x)$  is piecewise continuous
- (ii)  $\tilde{A}$  is normal if there exist  $x \in X$  such that  $\mu_{\tilde{A}}(x) = 1$
- (iii)  $\tilde{A}$  is convex iff  $\mu_{\tilde{A}}(\delta x_1 + (1 - \delta)x_2) \geq \min\{\mu_{\tilde{A}}(x_1), \mu_{\tilde{A}}(x_2)\}$  where  $\delta \in [0,1]$

### Triangular Fuzzy Number

The Fuzzy Number  $\tilde{A}^{TrFN} = (h_1, h_2, h_3)$  is a Triangular Fuzzy Number if its membership function is of the form

$$\mu_{\tilde{H}^{TrFN}}(x) = \begin{cases} 0 & x < h_1 \\ \frac{x - h_1}{h_2 - h_1} & h_1 < x < h_2 \\ 1 & x = h_2 \\ \frac{h_3 - x}{h_3 - h_2} & h_2 < x < h_3 \end{cases}$$

Where  $h_1 < h_2 < h_3$

### Ranking of Triangular Fuzzy Number

Let  $\tilde{H}^{TrFN} = (h_1, h_2, h_3)$  be a Triangular Fuzzy Number, by taking the co-efficient of fuzzy number from pascal's triangles and applying simple probability approach the following ranking formula were obtained

$$R(\tilde{H}^{TrFN}) = \frac{h_1 + 2h_2 + h_3}{4}$$

### Arithmetic Operations:

Let  $\tilde{H}^{TrFN} = (h_1, h_2, h_3)$  and Number  $\tilde{K}^{TrFN} = (k_1, k_2, k_3)$  be any two triangular fuzzy number then

$$\tilde{H}^{TrFN} + \tilde{K}^{TrFN} = (h_1 + k_1, h_2 + k_2, h_3 + k_3),$$

$$\tilde{H}^{TrFN} - \tilde{K}^{TrFN} = (h_1 - k_1, h_2 - k_2, h_3 - k_3)$$

### Trapezoidal Fuzzy Number

The Fuzzy Number  $\tilde{H}^{TraFN} = (h_1, h_2, h_3, h_4)$  is a Trapezoidal Fuzzy Number if its membership function is of the form

$$\mu_{\tilde{H}^{TraFN}}(x) = \begin{cases} 0 & x < h_1 \\ \frac{x - h_1}{h_2 - h_1} & h_1 < x < h_2 \\ 1 & h_2 < x < h_3 \\ \frac{h_4 - x}{h_4 - h_3} & h_3 < x < h_4 \\ 0 & x > h_4 \end{cases}$$

Where  $h_1 < h_2 < h_3 < h_4$

### Arithmetic Operations:

Let  $\tilde{H}^{TraFN} = (h_1, h_2, h_3, h_4)$  and  $\tilde{K}^{TraFN} = (k_1, k_2, k_3, k_4)$  be any two trapezoidal fuzzy number then

$$\tilde{H}^{TraFN} + \tilde{K}^{TraFN} = (h_1 + k_1, h_2 + k_2, h_3 + k_3, h_4 + k_4),$$

$$\tilde{H}^{TraFN} - \tilde{K}^{TraFN} = (h_1 - k_1, h_2 - k_2, h_3 - k_3, h_4 - k_4)$$

### Pentagonal Fuzzy Number

The Fuzzy Number  $\tilde{H}^{PFN} = (h_1, h_2, h_3, h_4, h_5)$  is a Pentagonal Fuzzy Number if its membership function is of the form

$$\mu_{\tilde{H}^{PFN}}(x) = \begin{cases} 0 & x < h_1 \\ \frac{x - h_1}{h_2 - h_1} & h_1 < x < h_2 \\ \frac{x - h_2}{h_3 - h_2} & h_2 < x < h_3 \\ 1 & x = h_3 \\ \frac{h_4 - x}{h_4 - h_3} & h_3 < x < h_4 \\ \frac{h_5 - x}{h_5 - h_4} & h_4 < x < h_5 \\ 0 & x > h_5 \end{cases}$$

Where  $h_1 < h_2 < h_3 < h_4 < h_5$

### Remark :

In the pentagonal fuzzy number  $(h_1, h_2, h_3, h_4, h_5)$  if  $\frac{h_2 + h_3 + h_4}{3} = h$  then it can be reduced into triangular fuzzy number as  $(h_1, h, h_5)$ .

### Ranking of Pentagonal Fuzzy Number

Let  $\tilde{A}^{PFN} = (h_1, h_2, h_3, h_4, h_5)$  be a Pentagonal Fuzzy Number, by taking the co-efficient of fuzzy number from pascal's triangles and applying simple probability approach the following ranking formula were obtained

$$R(\tilde{A}^{PFN}) = \frac{h_1 + 4h_2 + 6h_3 + 4h_4 + h_5}{16}$$

### Arithmetic Operations:

Let  $\tilde{H}^{PFN} = (h_1, h_2, h_3, h_4, h_5)$  and  $\tilde{K}^{PFN} = (k_1, k_2, k_3, k_4, k_5)$  be any two Pentagonal fuzzy number then

$$\tilde{H}^{PFN} + \tilde{K}^{PFN} = (h_1 + k_1, h_2 + k_2, h_3 + k_3, h_4 + k_4, h_5 + k_5),$$


$$\tilde{H}^{PFN} - \tilde{K}^{PFN} = (h_1 - k_1, h_2 - k_2, h_3 - k_3, h_4 - k_4, h_5 - k_5)$$

### Heptagonal Fuzzy Number

The Fuzzy Number  $\tilde{H}^{HEPFN} = (h_1, h_2, h_3, h_4, h_5, h_6, h_7)$  is a Heptagonal Fuzzy Number if its membership function is of the form

$$\mu_{\tilde{H}}(x) = \begin{cases} 0 & \text{for } x < h_1 \\ \frac{1}{2} \left( \frac{x - h_1}{h_2 - h_1} \right) & \text{for } h_1 \leq x \leq h_2 \\ \frac{1}{2} & \text{for } h_2 \leq x \leq h_3 \\ \frac{1}{2} + \frac{1}{2} \left( \frac{x - h_3}{h_4 - h_3} \right) & \text{for } h_3 \leq x \leq h_4 \\ \frac{1}{2} + \frac{1}{2} \left( \frac{h_5 - x}{h_5 - h_4} \right) & \text{for } h_4 \leq x \leq h_5 \\ \frac{1}{2} & \text{for } h_5 \leq x \leq h_6 \\ \frac{1}{2} \left( \frac{h_7 - x}{h_7 - h_6} \right) & \text{for } h_6 \leq x \leq h_7 \\ 0 & \text{for } x \geq h_7 \end{cases}$$

Where  $h_1 < h_2 < h_3 < h_4 < h_5 < h_6 < h_7$



**Remark :**

In the Heptagonal fuzzy number  $(h_1, h_2, h_3, h_4, h_5, h_6, h_7)$  if  $\frac{h_3+h_4+h_5}{3} = h$  then it can be reduced into pentagonal fuzzy number as  $(h_1, h_2, h, h_6, h_7)$ .

**Ranking of Heptagonal Fuzzy Number**

Let  $\tilde{H}^{HEPFN} = (h_1, h_2, h_3, h_4, h_5, h_6, h_7)$  be a Heptagonal Fuzzy Number, by taking the coefficient of fuzzy number from Pascal's triangles and applying simple probability approach the following ranking formula were obtained

$$R(\tilde{H}^{HEPFN}) = \frac{h_1 + 6h_2 + 15h_3 + 20h_4 + 15h_5 + 6h_6 + h_7}{64}$$

**Arithmetic Operations:**

Let  $\tilde{H}^{HEPFN} = (h_1, h_2, h_3, h_4, h_5, h_6, h_7)$   
and  $\tilde{K}^{HEPFN} = (k_1, k_2, k_3, k_4, k_5, k_6, k_7)$   
be any two Heptagonal fuzzy number then

$$\begin{aligned}\tilde{H}^{HEPFN} + \tilde{K}^{HEPFN} &= (h_1 + k_1, h_2 + k_2, h_3 + k_3, h_4 + k_4, h_5 + k_5, h_6 + k_6, h_7 + k_7), \\ \tilde{H}^{HEPFN} - \tilde{K}^{HEPFN} &= (h_1 - k_1, h_2 - k_2, h_3 - k_3, h_4 - k_4, h_5 - k_5, h_6 - k_6, h_7 - k_7)\end{aligned}$$

**Practical Situations:**

Some of the real time situations where the idea of assignment had been used is

- (i) Assigning Sales man to sales territory
- (ii) Assigning vehicles to appropriate routes
- (iii) Scheduling teachers to classes etc..

**Mathematical Formulation of an Assignment Problem**

An Assignment problem can be stated in the form of  $n \times n$  cost matrix  $(c_{ij})$  of real numbers in the following table

Jobs Persons	1	2	3	.....	N
1	$c_{11}$	$c_{12}$	$c_{13}$		$c_{1N}$
2	$c_{21}$	$c_{22}$	$c_{23}$		$c_{2N}$
3					
.					
.					
N	$c_{N1}$	$c_{N2}$	$c_{N3}$		$c_{NN}$

Mathematically assignment problem can be stated as

$$\text{Minimize } Z = \sum_{i=1}^n \sum_{j=1}^n c_{ij} x_{ij}$$

Subject to

$$\sum_{j=1}^n x_{ij} = 1 \quad i = 1, 2, \dots, n$$

$$\sum_{i=1}^n x_{ij} = 1 \quad j = 1, 2, \dots, n$$

Where  $x_{ij}$  is the decision variable denoting the assignment of the person  $i$  to job  $j$ ,  $c_{ij}$  is the cost of assignment of  $j^{th}$  job to the  $i^{th}$  person. The objective is to minimize the total cost of assigning all the jobs to the available persons

#### Algorithm:

**Step 1:** Examine whether the number of rows are equal to the number of columns in the given assignment problem. If not then a dummy row or dummy column must be added with zero cost element.

**Step 2:** Identify the least cost in each row of the cost matrix, subtract this least element from each element in that row. Then there will be at least one zero in each row which is termed as row reduction matrix.

**Step 3:** In the row reduction matrix identify the least element in each column and subtract from the remaining elements of the corresponding column. Which is termed as column reduction matrix.

**Step 4:** Examine the row successively until a row with exactly one zero is found. Encircle the zero element as an assigned cell and cross out all other zero in its column. If there are more than one zero, then pass onto the next row. The same procedure is followed until all the rows have been examined.

**Step 5:** Repeat the same procedure for all the columns of the reduced cost matrix. If there is no single zero in any row or column of the reduced matrix, then arbitrarily choose a row or column having the minimum number of zeroes. Arbitrarily choose zero in the row or column and cross the remaining zeroes in that row or column.

Repeat the steps 4 and 5 until all zeroes are either assigned or crossed out.

**Step 6:** An optimal assignment was obtained by using the above steps. Otherwise go to the remaining steps.

**Step 7:** Draw the minimum number of horizontal or vertical lines through all the zeroes as follows

(i) Mark (✓) to those rows where no assignment has been made

(ii) Mark (✓) to those columns which have zeroes in the marked rows

(iii) Mark (✓) rows which have assignments in marked columns

(iv) The process may be repeated until no more rows or columns to be examined

(v) Draw straight lines through all unmarked rows and marked columns.

**Step 8:** If the minimum number of lines passing through all the zeroes is equal to

	A	B	C	D
1	(2, 5, 8, 10, 12, 15, 17)	(10, 12, 15, 16, 17, 19, 21)	(5, 7, 9, 11, 13, 15, 17)	(1, 4, 5, 6, 7, 8, 9)
2	(3, 5, 7, 11, 13, 15, 18)	(0, 2, 4, 7, 9, 10, 12)	(2, 5, 8, 10, 12, 15, 17)	(10, 12, 15, 16, 17, 19, 21)
3	(0, 2, 4, 7, 9, 10, 12)	(1, 4, 5, 6, 7, 8, 9)	(6, 8, 10, 12, 14, 16, 18)	(5, 7, 9, 11, 13, 15, 17)
4	(1, 4, 5, 6, 7, 8, 9)	(5, 7, 9, 11, 13, 15, 17)	(10, 12, 15, 16, 17, 19, 21)	(0, 2, 4, 7, 9, 10, 12)

**Solution:**

**Step 1:**

The given assignment problem is balanced, therefore the heptagonal fuzzy

the number of rows or columns, the optimum solution is attained by an arbitrary allocation in the positions of the zeroes not crossed in step 3. Otherwise we shall move to the next step.

**Step 9:** Rewrite the cost matrix as follows

12 subtract this elements from all the uncrossed elements and add the same at the point of intersection of these lines. Other elements crossed by the lines remain unchanged.

Repeat the step 4 and 5 until an optimum solution were obtained.

### Numerical Example

**Consider the following assignment problem. Assign four jobs to four machine. The parameters were given in terms of Heptagonal Fuzzy Numbers.**

cost had been reduced to a crisp number by using Pascal's Graded Technique

A	B	C	D
---	---	---	---

$$\begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 9.9 & 15.8 & 11 & 5.9 \\ 10.04 & 6.54 & 9.9 & 15.8 \\ 6.54 & 5.9 & 12 & 11 \\ 5.9 & 11 & 15.8 & 6.54 \end{bmatrix}$$

**Step 2:**

Row reduced matrix

A      B      C      D

$$\begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 4 & 9.9 & 5.1 & 0 \\ 3.5 & 0 & 3.36 & 9.26 \\ 0.64 & 0 & 6.1 & 5.1 \\ 0 & 5.1 & 9.9 & 0.64 \end{bmatrix}$$

**Step 3:**

Column reduced matrix obtained from the above matrix

A      B      C      D

$$\begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 4 & 9.9 & 1.74 & 0 \\ 3.5 & 0 & 0 & 9.26 \\ 0.64 & 0 & 2.74 & 5.1 \\ 0 & 5.1 & 6.54 & 0.64 \end{bmatrix}$$

Allocating zeroes for each row and column the resultant assignment were obtained as follows

A      B      C      D

$$\begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 4 & 9.9 & 1.74 & (0) \\ 3.5 & \times & (0) & 9.26 \\ 0.64 & (0) & 2.74 & 5.1 \\ (0) & 5.1 & 6.54 & 0.64 \end{bmatrix}$$

**Assignment**  $1 \rightarrow D, 2 \rightarrow C, 3 \rightarrow B, 4 \rightarrow A$

**Total Assignment Value = 27.6 units**  
3413

(ii) In the above numerical example if  $\frac{h_3+h_4+h_5}{3} = h$  then it can be reduced into pentagonal fuzzy number as  $(h_1, h_2, h, h_6, h_7)$ .

Proceeding like the above example with Hungarian Method the

**Assignment is**  $1 \rightarrow D, 2 \rightarrow C, 3 \rightarrow B, 4 \rightarrow A$

**Total Assignment Value = 27.51 units**

**Comparison Table**

Fuzzy Numbers		Assignment Value
Heptagonal Fuzzy Numbers		27.60
Pentagonal Fuzzy Numbers		27.51

**Conclusion :**

In this paper the assignment problem were solved by Hungarian Method by using heptagonal fuzzy number ,further it had been converted into pentagonal fuzzy number .The results yield for both the fuzzy numbers represent the slight deviation.

**Reference**

- [1] R. Srinivasan , N. Karthikeyan ,A. Jayaraja ,A Proposed Ranking Method to solve Transportation Problem by Pentagonal Fuzzy Numbers ,Turkish



- Online Journal of Qualitative Inquiry, vol.12, Issue 3,
- [2] Shweta Singh ,G.C. Dubey, Rajesh Shrivastava ,A Comparative Analysis of Assignment Problem ,ISOR Journal of Engineering ,ISSN:2250-3021,vol 2,Issue 8,pp 1- 15,Aug 2021.
- [3] R. Queen Mary D. Selvi ,Solving Fuzzy Assignment Problem Using Centroid Ranking Method ,International Journal of Mathematics and its Applications,ISSN:2347 – 1557,6(3),9-16, 2018.
- [4] A. Arokiamary and S.Nithya ,Solving Fuzzy Assignment Problem for Hexagonal Fuzzy Number Using Revised Ones Assignment Method ,International Journal of Mathematicsand its Application ,ISSN :2347 – 1557 ,vol 4,issue 4 (2016) ,407-414.
- [5] Dr.A.Sahaya Sudha and M.Revathy, “A New Ranking on Hexagonal Fuzzy Numbers”, International journal of fuzzy logic systems (IJFLS) vol.6, No.4, October 2016.
- [6] Dr.ASahaya Sudha and S.Karunambigai ,”Solving a Transportation Problem Using Heptagonal Fuzzy Numbers ,International Journal of Advanced Research in Science, Engineering and Technology Vol. 4, Issue 1 , January 2017
- [7] A. Virgin Raj and S. Karthik, “Application of Pentagonal fuzzy number in Neural Network”. International journal of mathematics and its application. Volume 4, Issue 4(2016), 149-154(special issue) ISSN: 2347-1557.