

Fourteen Degree-based Topological Indices on twelve Penicillin Class of Drugs with QSPR Analysis

C.Tamilarasi¹, F. Simon Raj²

¹Hindustan Institute of Technology and Science, Chennai, Tamilnadu, India;

Email: tamilbabu4864@gmail.com

²Hindustan Institute of Technology and Science, Chennai, Tamilnadu, India;

E mail: simon240906@gmail.com

Received: 2022 March 15; **Revised:** 2022 April 20; **Accepted:** 2022 May 10.

Abstract- This research article aims to establish the values of fourteen notable topological indices that are defined on some chemical structures belonging to the Penicillin class of drugs, which are a type of antibiotic derived from Penicillin fungi. This is performed to predict the Physicochemical properties of these compounds through Quantitative Structure-Property Relationships (QSPR). Also, it is observed that these characteristics have a good correlation with the Physico-chemical characteristics of the selected chemical structures of the penicillin class of compounds. The chemical properties of the selected compounds that this paper discusses include Polarizability, Log Octanol-Water Partition Coefficient (Log K_{ow}), Molecular Volume, Molar Refractivity, Log P, Soil Absorption Coefficient (Koc), Log Koc.

MSC2010 codes 05C10, 05C30, 05C92, 05C90, 05C09

Keywords- Topological index, degree-based topological index, Quantitative Structure-Property Relationships, Penicillin, Physico-chemical properties.

1Introduction Chemical graph theory is one of the key branches of graph theory and plays a significant role in applied mathematics, which deals with chemical structures. Atoms represent vertices and chemical bonds represent edges of the

chemical structures in chemical graph theory. The numerical value associated to the vertices as well as edges in the molecular graph is termed as topological index/molecular descriptor. One of the main qualities of the topological indices is to predict the Physicochemical Properties of the respected chemical compounds through QSPR study^{2,9,10,12,13}. In this paper, some of the properties of the Penicillin class of drugs are taken for QSPR analysis with prominent fourteen degree-based topological indices which are given good correlation with them.

Penicillin was the first true antibiotic to be discovered, which helped to fight bacterial infections such as whooping cough, ear infection, Salmonella, and Tuberculosis. The Scottish bacteriologist Alexander Fleming first observed that colonies of the bacterium *Staphylococcus aureus* failed to grow in certain areas of its culture that had been accidentally contaminated by the mold *Penicillium notatum* which was green in color. This is the well-known *Penicillium notatum*, which was finally discovered in 1928.

Wiener index, the first topological index (distance-based) was introduced by Harold Wiener in 1947. It is defined as the summation of the shortest distance

between every single vertex of graph G . The progress of degree-based topological indices^{3,4} was started in 1975 by Milan Randić. Degree-based as well as distance-based topological indices are widely used in predicting the physicochemical properties^{7,11} of a class of compounds. Based on that, this paper is established. Here, fourteen notable degree-based topological indices are taken with twelve penicillin classes of drugs in the prediction task of properties through QSPR analysis.

2 Preliminaries In this part, some supportive notions are mentioned here, like, molecular graph G , degree of a vertex, topological index, degree-based topological indices, and basic formulas of fourteen degree-based topological indices. Let the molecular graph be the two-dimensional model of the chemical compound and is denoted as graph $G(V, E)$ holds two non-empty sets V (set of all atoms as vertices) & E (set of all chemical bonds as edges). The degree of vertex v is the number of vertices adjacent to the vertex v .

The basic definitions of the fourteen topological indices are as follows:

Definition 2.1 One of the important degree-based topological indices which have a highly predictive ability of Physico-chemical properties of chemical

compounds is the Augmented Zagreb Index ($Az(G)$). It was introduced by Furtula *et al* and is denoted as follows:

$$Az(G) = \sum_{u,v \in E} \left(\frac{dudv}{du+dv-2} \right)^3$$

Definition 2.2 Estrada *et al.* introduced the degree-based topological index named Atom Bond connectivity index ($ABC(G)$) which is a good predictor for the stability of alkanes and strain energy and is denoted as follows:

$$ABC(G) = \sum_{u,v \in E} \sqrt{\frac{du+dv-2}{dudv}}$$

Definition 2.3 The Geometric Arithmetic Index ($GA(G)$)¹ is the well-studied degree-based topological index and is denoted as follows:

$$GA(G) = \sum_{u,v \in E} \frac{2\sqrt{dudv}}{du+dv}$$

Definition 2.4 Harmonic Index ($H(G)$)¹⁵

$$H(G) = \sum_{u,v \in E} \frac{2}{du+dv}$$

Definition 2.5 Forgotten Index ($F(G)$)¹⁴ was defined to be used in the analysis of drug molecular structures which was introduced by B. Furtula *et al.* and is denoted as follows:

$$F(G) = \sum_{u,v \in E} du^2 + dv^2$$

Definition 2.6 The Zagreb indices $M_1(G)$, $M_2(G)$, and $M_3(G)$ are the numerical

measures to pronounce the structure-property relationships. $M_1(G)$ is the oldest degree-based topological index that was initiated in 1972 and the related versions were developed later.

$$M_1(G) = \sum_{u,v \in E} (du + dv)$$

$$M_2(G) = \sum_{u,v \in E} (du \, dv)$$

$$M_3(G) = \sum_{u,v \in E} |du - dv|$$

Definition 2.7 The Randic Index⁵ is one of the most powerful degree-based topological indices to study the structure-property relationships of graphs and is denoted as follows:

$$R(G) = \sum_{u,v \in E} \frac{1}{\sqrt{du \, dv}}$$

Definition 2.8 Sum Connectivity Index¹⁶ is denoted as follows:

$$S(G) = \sum_{u,v \in E} \frac{1}{\sqrt{du+dv}}$$

Definition 2.9 In 2013, Shirdel *et al.* proposed novel indices such as the First Hyper-Zagreb index ($H_1(G)$), Second Hyper-Zagreb index ($H_2(G)$) and are denoted as follows:

$$H_1(G) = \sum_{u,v \in E} (du + dv)^2$$

$$H_2(G) = \sum_{u,v \in E} (dudv)^2$$

Definition 2.10 FORAN index ($FR(G)$) holds the properties of the Forgotten index as well as the Randic index, and it is

suitable for predicting the relationship between molecular structures of drugs and their physicochemical properties. It is denoted as follows:

$$FR(G) = \sum_{u,v \in E} du \sqrt{\frac{du}{dv}} + dv \sqrt{\frac{dv}{du}}$$

Definition 2.11 Symmetric Division DegIndex($SDD(G)$)⁶ was defined as D.Vukicevic in 2010 by the order of its vertex degrees with the maximum degree and the minimum degree and is denoted as follows:

$$SDD(G) = \sum_{u,v \in E} \frac{\max(du, dv)}{\min(du, dv)} + \frac{\min(du, dv)}{\max(du, dv)}$$

Based on the above fourteen degree-based topological indices, we computed twelve penicillin classes of drugs. Table 1 and its continuation show the values of fourteen topological indices with the chemical structures of the penicillin class of compounds. Table 2 shows the results of the properties.

Figure 1 Molecular structures of Penicillin class of drugs




Table 1. It shows the calculated values of twelve Penicillin class of drugs with fourteen degree-based topological indices.

Penicillin class of drugs	$Az(G)$	$ABC(G)$	$GA(G)$	$H(G)$	$F(G)$	$M_1(G)$)	$M_2(G)$)
Amoxicillin	209.9397	19.6809	25.5873	10.885 7	390	138	168
Ampicillin	206.5647	18.8644	24.7615	10.585 7	370	132	161
Carbenicillin	221.3303	20.3475	26.6277	11.419 1	398	142	173
Dicloxacillin	276.1237	24.5956	32.3278	13.657 1	504	176	221
Nafcillin	274.1024	22.8761	30.7981	13.052 4	453	163	203
Oxacillin	239.522	22.3192	29.7213	12.552 5	440	158	195
Penicillin	195.6454	18.2454	18.0254	10.057 1	375	129	158
Ticarcillin	230.7237	21.1999	27.2766	11.523 7	448	158	190
Cloxacillin	267.3487	23.9189	31.3298	13.123 7	498	172	215
Methicillin	230.5803	21.5430	26.7407	11.552 5	398	142	175
Flucloxacillin	266.8772	23.8715	31.4533	13.219 0	480	170	211
Phenethicillin	267.1741	19.4049	25.7213	11.052 3	378	136	164

Table 1 continuation

Penicillin class of drugs	$M_3(G)$	$R(G)$	$S(G)$	$HM_1(G)$	$HM_2(G)$	$FR(G)$	$SDD(G)$
Amoxicillin	28	11.6308	11.9878	774	1326	159.0796	68.2500
Ampicillin	24	11.2370	11.6827	692	1177	150.0692	64.5822
Carbenicillin	26	12.1475	12.5909	744	1267	162.4651	69.9166
Dicloxacillin	34	14.5521	15.1634	946	1731	200.0976	84.5000
Nafcillin	25	13.6832	14.3792	859	1503	181.0681	76.0833
Oxacillin	26	13.2201	13.8936	830	1427	177.3061	74.9167
Penicillin	29	10.7570	11.1611	691	1212	150.9228	64.1666
Ticarcillin	32	12.4311	12.8607	828	1520	178.3217	75.6666
Cloxacillin	34	13.9867	14.6495	928	1671	197.7770	86.9998
Methicillin	26	12.2236	12.6400	748	1297	226.3317	68.5833
Flucloxacillin	30	14.0415	14.7101	902	1575	192.8531	81.5833
Phenethicillin	26	11.7201	12.1690	706	1168	126.4385	66.9167

Table 2. It shows the Penicillin class of drugs with their physicochemical properties

Penicillin class of drugs	Polarizability 10^{-24} cm^3	Log K_{ow}	Molar Volume	Molar Refractivity cm^3	Log P	K_{oc}	Log k_{oc}
Amoxicillin	36.3	0.97	236.2	91.5	0.61	865.5	115.7

n							
Ampicillin	35.7	1.45	239.3	89.9	1.35	534.4	105.4
Carbenicillin	36.7	1.19	246.2	92.6	1.01	2169	113.0
Dicloxacillin	44	3.86	290.1	111.0	3.02	9004	106.5
Nafcillin	43.7	3.79	289.9	110.3	3.52	1448	109.6
Oxacillin	40.2	2.57	268.5	101.3	2.05	3294	105.8
Penicillin	34.2	1.85	235.2	86.3	1.67	421.4	102.5
Ticarcillin	36	1.01	236.4	90.7	0.69	1176	117.3
Cloxacillin	42.1	3.22	279.3	106.2	2.53	5446	106.2
Methicillin	37.4	2.20	262.6	94.4	1.27	50.31	99.3
Flucloxacillin	42.1	3.42	283.7	106.3	2.60	9004	104.4
Phenethicillin	36.7	2.29	258	92.7	2.22	292.3	104.0

3 Results and discussions

Part 1 Here, the indices are correlated with the penicillin class of drugs and are

presented in table 3. High correlation coefficients are highlighted in the table and establish linear regression models for the highlighted values through QSPR analysis.

Table 3. It shows correlation coefficients between the physicochemical properties and indices.

Topological Index	Polarizability 10^{-24} cm^3	Log K_{ow}	Molar Volume	Molar Refractivity Cm^3	Log P	K_{oc}	Log K_{oc}
$A_z(G)$	0.8566	0.8404	0.8952	0.8593	0.8155	0.6169	0.4897
$ABC(G)$	0.9277	0.7935	0.8844	0.9271	0.6491	0.8135	0.6709
$GA(G)$	0.8885	0.6753	0.8268	0.8870	0.5719	0.7057	0.6488
$H(G)$	0.9694	0.8373	0.9242	0.9689	0.7335	0.8078	0.7167
$F(G)$	0.8664	0.7263	0.7783	0.8649	0.6091	0.8376	0.7799
$M_1(G)$	0.8959	0.7408	0.8129	0.8945	0.6294	0.8210	0.7708
$M_2(G)$	0.9201	0.7798	0.8430	0.9188	0.6663	0.8342	0.7670
$M_3(G)$	0.3654	0.3021	0.2720	0.3646	0.1785	0.6449	0.5569
$R(G)$	0.9558	0.8160	0.9023	0.9552	0.7055	0.8289	0.7370
$S(G)$	0.9623	0.8299	0.9110	0.9617	0.7266	0.8243	0.7421
$HM_1(G)$	0.8841	0.7133	0.7795	0.8824	0.5902	0.8320	0.7847
$HM_2(G)$	0.8264	0.6630	0.7121	0.8243	0.5353	0.8024	0.7524
$FR(G)$	0.5563	0.4655	0.5608	0.5557	0.2305	0.4579	0.1602
$SDD(G)$	0.8508	0.7016	0.7663	0.8495	0.5841	0.8378	0.7868

Part 2 In this section, suitable linear regression⁸ models (based on the highlighted values of table 3) of fourteen topological indices, and physicochemical properties of twelve penicillin class of drugs are established.

Linear regression models (QSPR analysis)

Linear regression model $N = n(\text{Topological index}) + n^\circ$

where N = physical property, TI = Topological index, n = slope & n° = intercept.

Using this, we obtained the following.

1. Augmented Zagreb Index

$$\text{Polarizability} = 0.1021(A_z(G)) + 14.212$$

$$\text{Log Kow} = 0.0309(A_z(G)) - 5.1145$$

$$\text{Molar Volume} = 0.6717(A_z(G)) + 98.9103$$

$$\text{Molar Refractivity} = 0.2591(Az(G)) + 35.4633$$

$$\text{Log P} = 0.0263(Az(G)) - 4.4479$$

2. Atom Bond Index

$$\text{Polarizability} = 1.4956(ABC(G)) + 6.7448$$

$$\text{Molar Volume} = 8.9797 (ABC(G)) + 68.2343$$

$$\text{Molar Refractivity} = 3.78171(ABC(G)) + 16.8166$$

$$\text{Koc} = 1244.774 (ABC(G)) - 23836.4$$

3. Geometric-Arithmetic Index

$$\text{Polarizability} = 0.7673(GA(G)) + 17.6341$$

$$\text{Molar Volume} = 4.4964 (GA(G)) + 136.6615$$

$$\text{Molar Refractivity} = 1.9381(GA(G)) + 44.4099$$

4. Harmonic Index

$$\text{Polarizability} = 0.7677(H(G)) + 17.6341$$

$$\text{Molar Volume} = 4.4964 (H(G)) + 136.6615$$

$$\text{Molar Refractivity} = 7.1243(H(G)) + 13.0583$$

$$\text{Koc} = 2228.133 (H(G)) - 23683.4$$

5. Forgotten Index

$$\text{Polarizability} = 0.0608(F(G)) + 12.7484$$

$$\text{Molar Refractivity} = 0.1536(F(G)) + 32.0659$$

$$\text{Koc} = 55.8028 (F(G)) - 21056.3$$

6. First Zagreb Index

$$\text{Polarizability} = 0.1847(M_1(G)) + 10.8079$$

$$\text{Molar Volume} = 1.0553 (M_1(G)) + 100.7412$$

$$\text{Molar Refractivity} = 0.4666(M_1(G)) + 27.1598$$

$$\text{Koc} = 160.6362 (M_1(G)) - 20500.9$$

7. Second Zagreb Index

$$\text{Polarizability} = 0.1406(M_2(G)) + 12.5793$$

$$\text{Molar Volume} = 0.8114 (M_2(G)) + 109.4039$$

$$\text{Molar Refractivity} = 0.3553(M_2(G)) + 31.6220$$

$$\text{Koc} = 120.9964 (M_2(G)) - 19716.7$$

8. Randic Index

$$\text{Polarizability} = 2.6713(R(G)) + 5.0046$$

$$\text{Log Kow} = 0.7039 (R(G)) - 6.5758$$

$$\text{Molar Volume} = 15.8801 (R(G)) + 59.7903$$

$$\text{Molar Refractivity} = 6.7561(R(G)) + 12.4161$$

$$\text{Koc} = 2198.565(R(G)) - 24972.1$$

9. Sum Connectivity Index

$$\text{Polarizability} = 2.6713(S(G)) + 5.0046$$

$$\text{Log Kow} = 0.6561 (S(G)) - 6.3137$$

$$\text{Molar Volume} = 14.694 (S(G)) + 67.111$$

$$\text{Molar Refractivity} = 6.2326(S(G)) + 15.763$$

$$\text{Koc} = 2003.8(S(G)) - 23556$$

10. First Hyper Zagreb Index

$$\text{Polarizability} = 0.0333(HM_1(G)) + 11.9924$$

$$\text{Molar Refractivity} = 0.0841(HM_1(G)) + 30.1714$$

$$\text{Koc} = 29.7338(HM_1(G)) - 21097.2$$

11. Second Hyper Zagreb Index

$$\text{Polarizability} = 0.0147(HM_2(G)) + 18.0735$$

$$\text{Molar Refractivity} = 0.0371(HM_2(G)) + 45.5608$$

$$\text{Koc} = 13.5559(HM_2(G)) - 16253.1$$

12. Symmetric Division Deg Index

$$\text{Polarizability} = 0.0147(HM_2(G)) + 18.0735$$

$$\text{Molar Refractivity} = 0.0371(HM_2(G)) + 45.5608$$

$$\text{Koc} = 13.5559(HM_2(G)) - 16253.1$$

The above twelve indices are highly correlated with some specific physicochemical properties of the Penicillin class of compounds (Polarizability, Log Octanol-Water Partition Coefficient (Log K_{ow}), Molecular Volume, Molar Refractivity, Log P, Soil Absorption Coefficient (Koc), Log Koc).

Hence, the QSPR analysis is done with the specific properties. The third Zagreb index and the FORAN index have moderately correlated properties. Geometric-Arithmetic index, Harmonic index, Randic index, and Sum Connectivity index have a very high correlation with the properties, especially with Polarizability and Molecular Refractivity that are underlined and highlighted in table 3. Figures 2 to 6 give a piece of solid information regarding this.




Figure 2 shows the correlation of $H(G)$ With Polarizability of Penicillin class of compounds.




Figure 3 shows the correlation of $H(G)$ With Molar Refractivity of Penicillin class of compounds.




Figure 4 shows the correlation of $R(G)$ With Polarizability of Penicillin class of compounds.




Figure 5 shows the correlation of $S(G)$ With Polarizability of Penicillin class of compounds.




Figure 6 shows the correlation of $S(G)$ With Molar Refractivity of Penicillin class of compounds.

Conclusion—In this paper, a QSPR analysis was established between fourteen degree-based topological indices and twelve

penicillin classes of drugs for the suitable pairs (topological index, property). In the field of pharmaceutical sciences, the properties of chemical compounds are very much required to establish new products. Instead of spending time, space for laboratory work, and the cost of research, topological indices are good alternators without these requirements. This theoretical approach obtained in this paper has promising aspects in designing new drugs.

References

1. K.C. Das, On geometric-arithmetic index of graphs, *MATCH Commun. Math. Comput. Chem*, 64 (2016), 619-630.
2. Devillers. A.T. Balaban, Topological indices and related descriptors in QSAR and QSPR, *God. & Bre. Sci. Pub., Netherland* (1999).
3. I. Gutman, B. Furtula, Three New/Old Vertex-Degree-Based Topological Indices, *MATCH Commun. Math. Comput. Chem.*, 72, (2014), 617-632.
4. Gutman, Degree-Based Topological Indices, *Croat. Chem. Acta*, 86, (4), (2013), 351-361.

5. Lucic, N. Trinajstić, Comparison between the sum-connectivity index and product-connectivity index for benzenoid hydrocarbons, *Che. Phy. Letters*, 475 **(2009)**, 146-148.
6. Marjan Matejić, Kinkar Ch. Das, Emina Milovanović, Bounds for Symmetric Division Deg Index of Graphs, *Filomat* 33:3 **(2019)**, 683-698.
7. Minati Kuanar, Ivan Gutman, Correlation of line graph parameters with physicochemical properties of octane isomers, *Indian Journal of Chemistry*, Vol. 38A **(1999)**, 525-528.
8. M. Randić, Comparative structure-property studies: Regressions using a single descriptor, *Croat. Chem. Acta* 66 **(1993)**.
9. Randić, Quantitative Structure-Property Relationship: boiling points and planar benzenoids, *New. J. Chem.* 20 **(1996)**, 1001-1009.
10. Sourav Mondal, Nilanjan De, QSPR analysis of some novel neighbourhood degree based topological descriptors, arXiv: 1906.06660v1 [cs.DM] **(2019)**.
11. Sunilkumar M. Hosamani, Correlation of domination parameters with physicochemical properties of octane isomers, 10.13140/RG.2.1.3808.4004 **(2015)**.
12. C. Tamilarasi, F. Simon Raj, QSPR Analysis of Novel Indices with Priority Polycyclic Aromatic Hydrocarbons (PAHs), *Tur. J. Com. & Math. Edu.*, Vol. 12 No. 10 **(2021)**, 3992-3999.
13. Vijayalaxmi Shigehalli, Rachanna Kanabur, QSPR Analysis of Degree-Based Topological Indices with physical properties of Benzenoid Hydrocarbons, *Gen. Lett. In Mat.*, Vol. 2, No. 3 **(2017)**, 150-169.
14. Wei Gao, Muhammad Kamran Siddiqui, Forgotten topological index of chemical structure in drugs, *Saudi Pharm. J.*, Vol. 24, No. 3 **(2016)**, 258-264.
15. L. Zhong, The harmonic index on graphs, *Appl. Math. Lett.* 25 **(2015)**, 561-566.
16. B. Zhou and N. Trinajstić, On a novel connectivity index, *J Math. Chem.* 46 **(2009)**, 1252-1270.